

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Sleep Health

Journal of the National Sleep Foundation

journal homepage: [sleephealthjournal.org](https://www.sleephealthjournal.org)

Goldilocks at work: Just the right amount of job demands may be needed for your sleep health

Monica E. Nelson, MA^{a,*}, Soomi Lee, PhD^a, Tammy D. Allen, PhD^b, Orfeu M. Buxton, PhD^c, David M. Almeida, PhD^d, Ross Andel, PhD^{e,f,g}

^a School of Aging Studies, University of South Florida, Tampa, Florida, USA

^b Department of Psychology, University of South Florida, Tampa, Florida, USA

^c Department of Biobehavioral Health, Pennsylvania State University, University Park, Pennsylvania, USA

^d Department of Human Development and Family Studies, Pennsylvania State University, University Park, Pennsylvania, USA

^e Center for Innovation in Healthy and Resilient Aging, Edson College of Nursing and Health Innovation, Arizona State University, Phoenix, Arizona, USA

^f Department of Neurology, Second Faculty of Medicine, Charles University and Motol University Hospital, Prague, Czech Republic

^g International Clinical Research Center, St. Anne's University Hospital Brno, Brno, Czech Republic

ARTICLE INFO

Article History:

Received 30 November 2021

Revised 29 July 2022

Accepted 3 September 2022

Available online xxx

Keywords:

Job demands

Job control

Job strain

Sleep

MIDUS

Non-linear relationships

ABSTRACT

Objectives: It has been reported that job demands affect sleep, but how different levels of job demands affect sleep remains unclear. We examined whether curvilinear relationships exist between job demands and multiple sleep health outcomes.

Design: Cross-sectional analyses with linear and quadratic effects, using self-administered survey data.

Setting: A national sample of US adults.

Participants: Workers from Midlife in the United States Study (MIDUS2; $n = 2927$).

Measurements: The Job Content Questionnaire assessed overall and 5 specific aspects of job demands (intensity, role conflict, work overload, time pressure, and interruptions). Habitual sleep health patterns across 5 dimensions (regularity, satisfaction/quality, daytime alertness, efficiency, and duration) were assessed. Age, sex, race/ethnicity, marital/partnered status, education, job tenure, work hours, body mass index, smoking status, and study sample were covariates.

Results: There were significant linear and quadratic relationships between job demands and sleep outcomes. Specifically, the linear effects indicated that participants with higher job demands had worse sleep health, such as shorter duration, greater irregularity, greater inefficiency, and more sleep dissatisfaction. The quadratic effects, however, indicated that sleep regularity and efficiency outcomes were the best when participants' job demands were moderate rather than too low or too high. These effects were found for overall job demands as well as for specific aspects of job demands. Stratified analyses further revealed that these curvilinear associations were mainly driven by participants with low job control.

Conclusions: Moderate levels of job demands, especially if combined with adequate job control, are related to optimal sleep health.

© 2022 The Authors. Published by Elsevier Inc. on behalf of National Sleep Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

Sleep is important to health, and poor sleep is related to negative health outcomes including cardiovascular disease,¹ dementia,² and mortality.¹ Therefore, identifying contributors to poor sleep is important to explain downstream health outcomes resulting from poor sleep. Sleep can be influenced by multiple factors at macro

(eg, environmental), social, and individual levels.³ Within the social level, work is important and has received much research attention in relation to sleep. Structural aspects of work, including long working hours⁴ and shift work,⁵ are negatively related to sleep outcomes. While many studies report the negative impact of work-related stressors on sleep,⁶ a complete lack of work-related stressors may also reflect poor engagement at work, which may have its own negative consequences.⁷ Similarly, the job demands-resources theory⁸ predicts that low job demands under certain conditions result in negative outcomes such as boredom⁹ and apathy. Additionally, certain types of work-related stressors appear to be more strongly related to negative sleep outcomes than others.¹⁰ Yet,

*Corresponding author: Monica Elizabeth Nelson, MA, University of South Florida, School of Aging Studies, 13301 Bruce B. Downs Blvd, MHC 1317, Tampa, FL 33612, USA. Tel.: 813-974-3232.

E-mail address: mnelson10@usf.edu (M.E. Nelson).

<https://doi.org/10.1016/j.sleh.2022.09.002>

2352-7218/© 2022 The Authors. Published by Elsevier Inc. on behalf of National Sleep Foundation. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Please cite this article as: M.E. Nelson et al., Goldilocks at work: Just the right amount of job demands may be needed for your sleep health, *Sleep Health* (2022), <https://doi.org/10.1016/j.sleh.2022.09.002>

most research examines only the linear relationship between work-related stressors and sleep (ie, more work is bad for sleep). This approach may provide biased results if there exists a range of work-related stressors that best promote health, including sleep, in working populations.

There is a theoretical basis for the possibility that some stress at work may facilitate health outcomes. For example, the hormetic stress concept,¹¹ motivated by the Yerkes-Dodson Law,¹² indicates that moderate, relative to low or high, levels of stressor exposure may relate to optimal performance or better health outcomes. Additionally, an ideal amount of stressors may benefit health.¹³ This may apply to sleep health. Much research on work-related stressors is guided by the job demands-control model,¹⁴ which posits that aspects of work can be categorized as job demands or as control resources. According to this model, many work characteristics (eg, long work hours, flexible hours, job insecurity, salary) are closely related to either job demands or to control, and the combination of high demands with low control results in job strain.¹⁴ Indeed, job demands or job strain (ie, work-related stressors) are the strongest predictors of employee health and well-being outcomes.^{15,16} Recent systematic reviews show that high job demands and job strain are related to sleep disturbances¹⁷ and poor sleep quality.¹⁸ Both demands and strain have evidenced negative linear relationships with sleep outcomes in working adults from Europe¹⁹ and Asia.^{20,21} Yet, there may be untested non-linear relationships, such that the levels of sleep health are lower when job demands are too little or too high. For example, research on job autonomy and well-being illustrates that both too much and too little autonomy may relate to poorer well-being.²² Regarding non-linear relationships between job demands and sleep outcomes, one study found non-significant quadratic effects for the dimensions of the job demands-control-support model,²³ but assessed only sleep quality, limiting conclusions regarding other aspects of sleep health. Moreover, most studies assessed job demands broadly, preventing examination of whether specific aspects of job demands may relate to sleep health outcomes both linearly and non-linearly.

The current study

Motivated by the hormetic stress concept,¹¹ we aimed to assess linear and curvilinear relationships between job demands and sleep health outcomes using a national sample of workers. In examining these relationships, we used an overall job demands composite as well as the individual items of the job demands scale. We also assessed 5 dimensions of sleep health (ie, regularity, satisfaction, alertness, efficiency, and duration²⁴). We expected that moderate job demands would be associated with better sleep outcomes, with both low and high job demands associated with worse sleep outcomes. Additionally, we examined differences by the level of job control to tap into the job strain concept¹⁴ (ie, high job demands, low job control).

Participants and methods

Data source and participants

We used data from the National Survey of Midlife in the United States (MIDUS; www.midus.wisc.edu), which assesses behavioral, psychological, and social factors related to variation in mental and physical health in the process of aging.²⁵ Specifically, data were from the first longitudinal follow-up (MIDUS2; in 2004–2006) and the refresher sample (MIDUS Refresher; in 2011–2014). Data from the Milwaukee, Wisconsin subsamples that collected information on Black/African American participants were included in the MIDUS2 and MIDUS Refresher samples. All variables were self-reported. Of the combined MIDUS2 and MIDUS Refresher samples ($n = 9640$),

participants were excluded if they (1) did not complete the self-administered questionnaire (SAQ) that included job demands questions ($n = 2278$), (2) were not currently paid workers ($n = 3576$), (3) did not work at least 30 hours per week ($n = 735$), and (4) did not have valid data for the job demands composite question ($n = 124$). Thus, the final analytic sample was 2927 workers. The MIDUS study was approved by all appropriate Institutional Review Boards, and all MIDUS participants provided informed consent. The current study used publicly available, de-identified data and was exempt from local IRB review.

Measures

Sleep health

Five dimensions of sleep health were included as the dependent variables. These dimensions follow the Ru-SATED model,²⁴ which includes sleep regularity, satisfaction, alertness, timing, efficiency, and duration. Sleep timing was not collected in the MIDUS survey; therefore, we did not include this dimension. In the current study, we refer to the available sleep variables as irregularity, dissatisfaction, nap frequency (a proxy for daytime alertness), inefficiency, and duration, respectively.

Sleep irregularity was measured as the absolute value of the difference between weekday and weekend sleep duration; higher scores indicated more irregular sleep. *Sleep dissatisfaction* ($\alpha = .80$) was measured as the mean of 4 items (feeling unrested during the day; trouble falling asleep; waking up during the night and having difficulty going back to sleep; waking up too early in the morning and being unable to go back to sleep) assessed on a 1–5 scale; higher scores indicated more dissatisfying sleep. *Nap frequency* was measured through a question indicating the number of naps that lasted 5+ minutes participants would take during a usual week; higher scores indicated more frequent naps (ie, less daytime alertness). We assessed nap frequency as it was the only available survey measure related to daytime alertness. In the original conceptualization of sleep health,²⁴ alertness, sleepiness, and napping were grouped, because these concepts are all related to levels of homeostatic sleep pressure.²⁶ Homeostatic sleep pressure accumulates during the day with continued wakefulness.²⁶ Sleep propensity and sleepiness are driven by this homeostatic process and increase the likelihood of napping.²⁷ Napping subsequently reduces homeostatic sleep pressure²⁸ and increases alertness and vigilance.²⁹ Yet, napping has been related to several negative outcomes,²⁷ including cardiovascular events,^{30,31} Alzheimer's disease,³² and mortality,³⁰ potentially because it is often accompanied by poor nighttime sleep.³¹ As a result, napping has garnered evidence for its utility as a sleep health measure²⁴ and a proxy for less daytime alertness. *Sleep inefficiency* was measured by sleep onset latency with a question assessing the time it takes participants to fall asleep at bedtime; higher scores indicated more inefficient sleep (in hours). While some previous research used wake after sleep onset (WASO) together with sleep onset latency,³³ we chose to use sleep onset latency alone as actigraphy-measured WASO was only available in a sub-sample of MIDUS. Sleep onset latency has been frequently used to indicate sleep inefficiency.³⁴ Finally, *sleep duration* was measured with a question asking how much sleep participants usually get at night on weekdays (in hours).

Job demands

Job demands indicators, the independent variables, were assessed using the Job Content Questionnaire (JCQ³⁵). Specifically, as suggested previously,³⁵ the following 5 items from the Demands Scale of the JCQ³⁵ were used as individual indicators of job demands, and the composite of the 5 items was used as the indicator for overall

demands: Participants reported how often (1 = all of the time, 2 = most of the time, 3 = some of the time, 4 = little of the time, 5 = never) the following aspects of their jobs occurred, on average or in the past year: (1) *intensity* (“do you have to work very intensively?”), (2) *role conflict* (“do different people or groups at work demand things from you that you think are hard to combine?”), (3) *work overload* (“do you have too many demands made on you?”), (4) *time pressure* (“do you have enough time to get everything done?”), and (5) *interruptions* (“do you have lots of interruptions?”). Intensity, role conflict, work overload, and interruptions were reverse coded such that higher scores indicated more demands. The *overall demands* item summed the 5 job demands questions with higher scores indicating higher demands ($\alpha = .74$).

Job control

Job control was assessed using the JCQ³⁵-based scales for skill discretion and decision authority, plus an additional question separate from the JCQ that asked participants about their perception of overall job control. The skill discretion score summed 3 items measuring how often on average (1) participants learned new things at work, (2) participants' work demanded a high level of skill or expertise, and (3) their job provided them with a variety of things that interested them. The decision authority score summed 6 items measuring how often participants had (1) to initiate things, (2) a choice in deciding how to do work tasks, (3) a choice in deciding what tasks to do, (4) a say in decisions about their work, (5) a say in planning their work environment, and how often participants could control the amount of time they spent on tasks. The one question that was separate from the JCQ asked how much control participants thought they had over their work situation (0 = no control at all to 10 = very much control). The 2 scale scores and the single item were first standardized with z-scores then averaged to create the *job control* factor ($\alpha = .67$). Job control was then dichotomized (low job control < median; high job control \geq median), following prior research.³⁶

Covariates

We considered sociodemographic and health-related covariates that may be associated with work experiences and sleep. Sociodemographic variables included age (years), sex (female = 0; male = 1), race/ethnicity (non-Hispanic White = 0; Other race/ethnicity = 1), marital/partnered status (unpartnered [ie, separated, divorced, widowed, never married, not living with partner] = 0; married/cohabitating = 1), final education level (1 = no school to 12 = professional degrees), job tenure (years), and average work hours. Health-related variables included body mass index and current smoking status (no = 0; yes = 1). Additionally, to account for potential sample differences, we adjusted for the sample identifier (MIDUS2 = 0; MIDUS Refresher = 1). Continuous covariates were centered at the sample means.

Statistical analyses

First, we assessed our main study variables for normality. Results indicated that variables were within an acceptable range, based on previous research with other continuous health variables^{37,38} (ie, < 3 for skewness and < 10 for kurtosis³⁹), to proceed with analyses assuming normality. Second, we conducted descriptive statistics and intercorrelations of study variables. Next, separate hierarchical regression analyses were conducted to assess the relationship between the various job demands indicators and each of the 5 sleep health dimensions. The 6 job demands indicators (overall composite + 5 individual items) were centered at the sample mean for analyses. Quadratic effects of the job demands indicators were created by multiplying each of the centered job demands items by itself (eg,

intensity x intensity). Inclusion of quadratic effects improved model fit when significant quadratic effects were found, as indicated by smaller RMSE values and larger adjusted R^2 values. Adding cubic effects did not significantly improve model fit—thus, cubic effects were not included in the models. Unadjusted models assessed the linear and quadratic effects of each job demands indicator on sleep health. Adjusted models included covariate effects (Model 1), linear effects of job demands (Model 2), and quadratic effects of job demands (Model 3). Missing data were excluded through listwise deletion for each analysis. To tap into the job strain concept,¹⁴ we also conducted analyses stratified by level of job control (high and low) to assess whether the effects of demands may depend on job resources available to participants. Finally, dose-response relationships were assessed between job demands and sleep health for all significant quadratic effects. All analyses were conducted with SAS Version 9.4, with significance assessed at $p < .05$ (2-tailed).

Results

Table 1 includes descriptive statistics of study variables. Participants were approximately 48 years old on average ($SD = 10.04$, $Range = 24-81$), with slightly more than half the sample being male (51.62%), and a majority of the sample being non-Hispanic White (82.32%). The sample was well-educated (≥ 4 -year college education = 48.39%; \geq high school diploma or equivalent = 96.96%). Most of the sample were either married or cohabitating (72.56%) and were not current smokers (86.16%).

Supplemental Table 1 includes correlations among the main study variables. Job control was weakly related to job demands indicators ($r_s = .06-.15$). The general associations between job demands indicators and sleep health dimensions indicated higher demands were typically related to poorer sleep health, though not all correlations were significant. One exception was found for nap frequency, in which higher job demands were related to lower nap frequency, but the magnitude of the correlation was very small ($r = -.05$). Higher job control was related to better sleep health. Correlations among sleep health dimensions were in the expected directions; yet, the relationships were weak overall ($r_s < |.25|$) except a moderate correlation between sleep inefficiency and sleep dissatisfaction ($r = .45$), indicating that each dimension conveys unique information about sleep health.

Linear and quadratic relationships between job demands and sleep health

Table 2 includes results from unadjusted and adjusted models for sleep duration. Controlling for study sample, age, sex, race/ethnicity, marital/partnered status, education, job tenure, work hours, body mass index, and smoking status, significant linear associations were found for 2 of the job demands indicators. Specifically, higher overall job demands ($b = -0.02$, $SE = 0.01$, $p = .014$) and intensity ($b = -0.09$, $SE = 0.03$, $p = .002$) were related to shorter sleep duration. No significant quadratic effects were found in fully adjusted models ($p_s \geq .054$).

Table 3 includes results from adjusted models for sleep irregularity, sleep inefficiency, nap frequency, and sleep dissatisfaction (unadjusted effects in Supplemental Table 2). Beginning with sleep irregularity, in fully adjusted models, there were linear associations between each of the 6 job demands indicators and sleep irregularity, such that higher job demands overall and each domain (intensity, role conflict, work overload, time pressure, and interruptions) were related to more irregular sleep. There were also significant quadratic effects of overall demands and work overload on sleep irregularity (**Fig. 1A** and **B**). Sleep was most regular around moderate levels of both overall demands and work overload.

Next, results for sleep inefficiency revealed significant linear relationships, such that higher overall demands ($b = 0.005$, $SE = 0.002$,

Table 1
Descriptive statistics for study variables

Variable	N	Missing	M/N	SD/%
Sociodemographic and health variables				
Sample	2927	0		
MIDUS2			1847	63.10%
MIDUS Refresher			1080	36.90%
Age, y	2927	0	48.03	10.04
Sex	2927	0		
Male			1511	51.62%
Female			1416	48.38%
Race/Ethnicity	2919	8		
Non-Hispanic White			2403	82.32%
Other			516	17.68%
Marital/partnered status	2923	4		
Not married or partnered ^a			802	27.44%
Married or cohabitating			2121	72.56%
Education ^b	2924	3		
Less than high school			89	3.04%
High school graduate or equivalent			603	20.62%
Some college or 2-y degree			817	27.94%
College graduate (4+ y) or higher			1415	48.39%
Job tenure, y	2877	50	24.80	11.16
Work hours	2927	0	43.28	8.02
Body mass index	2800	127	28.37	6.16
Current smoker	2927	0		
No			2522	86.16%
Yes			405	13.84%
Job demands variables				
Job demands composite	2927	0	15.41	3.12
Intensity	2919	8	3.72	0.77
Role conflict	2917	10	2.67	0.91
Work overload	2926	1	2.89	0.87
Time pressure	2923	4	2.63	0.98
Interruptions	2923	4	3.50	0.93
Job control variables				
Job control	2917	10	6.67	2.41
Skill discretion composite	2921	6	10.67	2.18
Decision authority composite	2921	6	21.80	4.39
Job control composite	2927	0	0.00	0.78
Sleep health variables				
Sleep duration	2915	12	6.82	1.09
Sleep irregularity	2866	61	1.02	1.12
Sleep inefficiency	2889	38	0.39	0.37
Nap frequency	2900	27	1.47	2.16
Sleep dissatisfaction	2917	10	2.46	0.82

^a Includes separated, divorced, widowed, never married, not living with partner.

^b There are 12 educational categories in MIDUS that we have grouped into the 4 categories.

Note. Abbreviations: M = mean. SD = standard deviation. Intensity = work intensively at job. Role Conflict = work demands hard to combine. Work Overload = too many demands at job. Time Pressure = time to get everything done at job. Interruptions = lot of interruptions at job. Higher scores indicate higher demands.

$p = .048$), intensity ($b = 0.02$, $SE = 0.01$, $p = .011$), and interruptions ($b = 0.02$, $SE = 0.01$, $p = .048$) were related to more inefficient sleep. A significant quadratic relationship appeared between work overload and sleep inefficiency ($b = 0.02$, $SE = 0.01$, $p < .001$). Fig. 1C illustrates that sleep inefficiency was lowest around moderate levels of work overload. Results for nap frequency revealed a significant linear association with intensity ($b = -0.11$, $SE = 0.05$, $p = .042$), with more intense work associated with less frequent naps (ie, more alertness). There were no significant quadratic relationships between any of the job demands indicators and nap frequency ($ps \geq .067$).

Finally, fully adjusted models for sleep dissatisfaction revealed linear relationships between all 6 job demands indicators and sleep dissatisfaction. Higher overall demands ($b = 0.04$, $SE = 0.01$, $p < .001$), intensity ($b = 0.06$, $SE = 0.02$, $p = .003$), role conflict ($b = 0.10$, $SE = 0.02$, $p < .001$), work overload ($b = 0.16$, $SE = 0.02$, $p < .001$), time pressure ($b = 0.08$, $SE = 0.02$, $p < .001$), and interruptions ($b = 0.09$, $SE = 0.02$, $p < .001$) were related to more dissatisfied sleep. No significant quadratic relationships appeared between any of the job demands indicators and sleep dissatisfaction ($ps \geq .102$).

Table 2
Linear and quadratic associations between job demands and sleep duration

Variable	Unadjusted sleep duration			Adjusted sleep duration		
	b	SE	p	b	SE	p
Overall demands	-0.01	0.01	.032	-0.02	0.01	.014
Overall demands ²	-0.002	0.001	.259	-0.001	0.001	.652
Intensity	-0.09	0.03	<.001	-0.09	0.03	.002
Intensity ²	-0.01	0.02	.662	0.002	0.02	.927
Role conflict	0.01	0.02	.689	-0.003	0.02	.906
Role conflict ²	-0.05	0.02	.003	-0.04	0.02	.054
Work overload	-0.03	0.02	.203	-0.03	0.02	.220
Work overload ²	-0.05	0.02	.005	-0.03	0.02	.074
Time pressure	-0.01	0.02	.603	-0.03	0.02	.274
Time pressure ²	-0.03	0.02	.090	-0.01	0.02	.431
Interruptions	-0.03	0.02	.116	-0.04	0.02	.050
Interruptions ²	-0.03	0.02	.172	-0.01	0.02	.604

Note. Overall job demands and specific items were entered in separate models. Bolded values indicate significant effects. Abbreviations: b = unstandardized regression estimate. SE = standard error. Overall Demands = job demands composite score ($n = 2915$ unadjusted; $n = 2734$ adjusted). Intensity = work intensively at job ($n = 2908$ unadjusted; $n = 2729$ adjusted). Role Conflict = work demands hard to combine ($n = 2905$ unadjusted; $n = 2727$ adjusted). Work Overload = too many demands at job ($n = 2914$ unadjusted; $n = 2733$ adjusted). Time Pressure = time to get everything done at job ($n = 2912$ unadjusted; $n = 2732$ adjusted). Interruptions = lot of interruptions at job ($n = 2911$ unadjusted; $n = 2731$ adjusted). Adjusted models controlled for study sample, age, sex, race/ethnicity, marital status, education, job tenure, work hours, body mass index, and whether the participant currently smoked. Continuous covariates and main predictors were centered at the sample mean. Bolded values indicate significant effects at $p < .05$.

Table 3
Linear and quadratic associations between job demands and sleep health dimensions

Variable	Sleep irregularity			Nap frequency		
	b	SE	p	b	SE	p
Overall demands	0.03	0.01	<.001	-0.01	0.01	.548
Overall demands ²	0.003	0.002	.031	-0.004	0.003	.187
Intensity	0.08	0.03	.006	-0.11	0.05	.042
Intensity ²	0.03	0.03	.185	-0.09	0.05	.067
Role conflict	0.06	0.03	.018	0.07	0.05	.137
Role conflict ²	0.03	0.02	.110	-0.03	0.04	.371
Work overload	0.06	0.03	.017	0.03	0.05	.605
Work overload ²	0.04	0.02	.039	-0.06	0.04	.145
Time pressure	0.05	0.02	.027	-0.04	0.05	.426
Time pressure ²	0.02	0.02	.316	-0.04	0.04	.285
Interruptions	0.07	0.02	.003	-0.05	0.05	.287
Interruptions ²	0.04	0.02	.078	-0.04	0.04	.251
<hr/>						
	Sleep inefficiency			Sleep dissatisfaction		
	b	SE	p	b	SE	p
Overall demands	0.005	0.002	.048	0.04	0.01	<.001
Overall demands ²	0.001	0.001	.246	-0.0003	0.001	.775
Intensity	0.02	0.01	.011	0.06	0.02	.003
Intensity ²	0.001	0.01	.920	-0.01	0.02	.623
Role conflict	0.01	0.01	.505	0.10	0.02	<.001
Role conflict ²	0.01	0.01	.124	-0.01	0.01	.275
Work overload	0.02	0.01	.054	0.16	0.02	<.001
Work overload ²	0.02	0.01	<.001	-0.01	0.01	.695
Time pressure	-0.003	0.01	.725	0.08	0.02	<.001
Time pressure ²	0.002	0.01	.781	-0.02	0.01	.110
Interruptions	0.02	0.01	.048	0.09	0.02	<.001
Interruptions ²	-0.005	0.01	.461	-0.02	0.01	.102

Note. Sample sizes for analyses ranged from $n = 2685$ to $n = 2737$. Overall job demands and specific items were entered in separate models. Bolded values indicate significant effects. Abbreviations: b = unstandardized regression estimate. SE = standard error. Overall Demands = job demands composite score. Intensity = work intensively at job. Role Conflict = work demands hard to combine. Work Overload = too many demands at job. Time Pressure = time to get everything done at job. Interruptions = lot of interruptions at job. Models controlled for study sample, age, sex, race/ethnicity, marital status, education, job tenure, work hours, body mass index, and whether the participant currently smoked. Continuous covariates and main predictors were centered at the sample mean. Bolded values indicate significant effects at $p < .05$.




Fig. 1. Curvilinear relationships between job demands indicators and sleep health dimensions. Overall job demands ($M = 15.41$) and work overload ($M = 2.89$) were mean centered for analyses. Work overload represents the frequency (1 = never, 2 = little of the time, 3 = some of the time, 4 = most of the time, 5 = all of the time) with which participants had too many demands made on them. Overall job demands represents the composite of the 5 job demands indicators (range = 5–25). Higher values on sleep health variables represent poorer sleep. All figures include bar graphs with frequency (%) of responses for sleep health (right) and job demands (top). A scatterplot represents the responses participants provided. Shaded regions represent 95% confidence intervals. A) A significant quadratic relationship exists between overall job demands and sleep irregularity ($b = 0.003$, $SE = 0.002$, $p < .001$), such that moderate demands were associated with more regular sleep. B) A significant quadratic relationship exists between work overload and sleep irregularity ($b = 0.04$, $SE = 0.02$, $p = .039$), such that moderate demands were associated with more regular sleep. C) A significant quadratic relationship exists between work overload and sleep inefficiency ($b = 0.02$, $SE = 0.01$, $p < .001$), such that moderate demands were associated with more efficient sleep.

Associations between job demands and sleep health stratified by job control

Table 4 includes results from unadjusted and adjusted models for overall demands and sleep health for participants with low and high job control. In fully adjusted models, overall demands were linearly associated with sleep inefficiency ($b = 0.01$, $SE = 0.003$, $p < .001$) only for participants with low job control. Overall demands were related to greater sleep irregularity (low control: $b = 0.03$, $SE = 0.01$, $p < .001$; high control: $b = 0.02$, $SE = 0.01$, $p = .048$) and greater sleep dissatisfaction (low and high control: $b = 0.04$, $SE = 0.01$, $p < .001$) in both participants with low and high job control. No significant quadratic effects of overall demands appeared for sleep health stratified by job control ($ps \geq .079$).

Supplemental Tables 3–7 include results for linear and quadratic relationships between each of the specific job demands indicators and each sleep health outcome stratified by job control. For the sake of parsimony, results consistently found between unadjusted and adjusted models are described here. There were nuanced differences by models, yet overall, the negative linear relationship between job demands and sleep was more apparent for individuals with low job control. The quadratic relationship was consistently found for specific job demands indicators, such that moderate job demands were associated with better sleep health (ie, longest sleep duration, more regular sleep, and more efficient sleep; Fig. 2) mostly for individuals with low job control. A notable exception was observed in sleep dissatisfaction. Each of the 5 job demands indicators were linearly related to sleep dissatisfaction, such that higher demands

Table 4
Linear and quadratic associations between overall demands and sleep characteristics

Variable	Unadjusted effects						Covariate adjusted effects					
	Low control			High control			Low control			High control		
	b	SE	p	b	SE	p	b	SE	p	b	SE	p
<i>Sleep duration</i> ^a												
Overall demands	-0.02	0.01	.047	-0.01	0.01	.162	-0.02	0.01	.062	-0.02	0.01	.072
Overall demands ²	-0.003	0.002	.103	0.0004	0.002	.847	-0.003	0.002	.198	0.002	0.002	.441
<i>Sleep irregularity</i> ^b												
Overall demands	0.03	0.01	<.001	0.03	0.01	.007	0.03	0.01	<.001	0.02	0.01	.048
Overall demands ²	0.004	0.002	.064	0.005	0.002	.029	0.004	0.002	.079	0.003	0.002	.127
<i>Sleep inefficiency</i> ^c												
Overall demands	0.01	0.003	.015	-0.0005	0.003	.884	0.01	0.003	.010	0.0002	0.003	.963
Overall demands ²	0.001	0.001	.077	0.001	0.001	.114	0.001	0.001	.431	0.001	0.001	.269
<i>Nap frequency</i> ^d												
Overall demands	-0.02	0.02	.378	-0.05	0.02	.009	-0.01	0.02	.790	-0.01	0.02	.492
Overall demands ²	-0.005	0.004	.245	-0.001	0.004	.821	-0.01	0.004	.112	-0.001	0.004	.789
<i>Sleep dissatisfaction</i> ^e												
Overall demands	0.04	0.01	<.001	0.04	0.01	<.001	0.04	0.01	<.001	0.04	0.01	<.001
Overall demands ²	0.0003	0.001	.842	0.0005	0.002	.774	-0.005	0.002	.767	-0.0004	0.002	.930

Bolded values indicate significant effects at $p < .05$.

^a Unadjusted low control ($n = 1458$), unadjusted high control ($n = 1457$), adjusted low control ($n = 1363$), adjusted high control ($n = 1371$).

^b Unadjusted low control ($n = 1433$), unadjusted high control ($n = 1433$), adjusted low control ($n = 1342$), adjusted high control ($n = 1349$).

^c Unadjusted low control ($n = 1440$), unadjusted high control ($n = 1449$), adjusted low control ($n = 1345$), adjusted high control ($n = 1364$).

^d Unadjusted low control ($n = 1453$), unadjusted high control ($n = 1447$), adjusted low control ($n = 1358$), adjusted high control ($n = 1362$).

^e Unadjusted low control ($n = 1460$), unadjusted high control ($n = 1457$), adjusted low control ($n = 1365$), adjusted high control ($n = 1372$).

Note. Bolded values indicate significant effects. Adjusted models controlled for study sample, age, sex, race/ethnicity, marital status, education, job tenure, work hours, body mass index, and whether the participant currently smoked. Low control and high control were stratified based on a median split (\geq median = high, $<$ median = low). Continuous covariates and main predictors were centered at the sample mean. Abbreviations: b = unstandardized regression estimate. SE = standard error.




Fig. 2. Curvilinear relationships between work overload and sleep health dimensions stratified by job control. Work overload ($M = 2.89$) was mean centered for analyses. Work overload represents the frequency (1 = never, 2 = little of the time, 3 = some of the time, 4 = most of the time, 5 = all of the time) with which participants had too many demands made on them. Lower values on sleep duration represent poorer sleep (Panels A and B) whereas higher values on sleep irregularity and sleep inefficiency represent poorer sleep (Panels C-F). Analyses were stratified by job control (left: low control; right: high control). All figures include bar graphs with frequency (%) of responses for sleep health (right) and work overload (top). A scatterplot represents the responses participants provided. Shaded regions represent 95% confidence intervals. Significant quadratic relationships exist between work overload and sleep duration (A: low control), sleep irregularity (C: low control), and sleep inefficiency (E: low control; F: high control). Moderate demands were associated with longer sleep duration, and more regular and efficient sleep. Relationships between work overload, sleep duration, and sleep irregularity were non-significant for participants with high job control. There were no significant quadratic relationships between work overload and either nap frequency or sleep dissatisfaction. There were also no other significant quadratic effects of the remaining job demands indicators for any of the sleep health dimensions in stratified analyses.

were related to more dissatisfied sleep similarly for participants with low and high job control.

Dose-response relationships between job demands and sleep health

Dose-response relationships were assessed for all significant quadratic effects. For the association between overall demands and sleep irregularity, the most regular sleep was estimated for job demands around 14 out of 25. This can be translated as “moderate”, or having job demands “little” (= 2) to “some” (= 3) of the time across the 5 items. For the association between work overload and irregularity, the most regular sleep was estimated for job demands 2 out of 5. For the association between work overload and sleep inefficiency, the most efficient sleep was estimated for job demands corresponding to 3 out of 5.

For the stratified analyses (Fig. 2), the most optimal sleep health for participants with low job control was estimated at 2 out of 5 on work overload for both sleep duration (longest duration) and sleep irregularity (most regular sleep). Efficient sleep was estimated for work overload corresponding to 3 out of 5 for both participants with low and high job control.

Discussion

We found initial empirical evidence that not all work-related stressors may be harmful for sleep health, providing support for the use of the hormetic stress concept¹¹ in sleep research. We assessed linear and curvilinear relationships between job demands and sleep health. Consistent with prior research, we found a general pattern that higher job demands were related to poorer sleep health. However, for some sleep health dimensions, as hypothesized, we found a nonlinear pattern of relationships. Specifically, curvilinear relationships did exist between job demands and certain sleep health dimensions (ie, irregularity, inefficiency), with moderate demands related to more regular and efficient sleep. Results stratified by job control further revealed that these linear and quadratic relationships appeared stronger for participants with low job control, indicating a potential exacerbating effect of low job control. Evidence suggests that both too low—potentially reflecting disengagement at work—and too high job demands may be maladaptive⁷ to sleep health (consistent with the hormetic stress concept¹¹), indicating that a moderate amount of stressor exposure is associated with optimal sleep.

Overall, findings support the job demands-control model,¹⁴ which suggests job demands are related to different workplace and health outcomes that may be attenuated by level of job control. Instead of only examining whether the combination of high job demands with low job control was associated with worse sleep outcomes, we assessed whether certain levels of job demands were associated with sleep outcomes, based on the hormetic stress concept.¹¹ The application of this dose-response approach between job demands and sleep outcomes results in a more nuanced understanding of relationships between work-related stressors and health, suggesting that an ideal level of psychosocial stressors may be best for sleep health (c.f., eustress⁴⁰). Specifically, we found that exposure to moderate levels of job demands can be beneficial for sleep health.

Moderate job demands were associated with more optimal sleep regularity and efficiency. While high job demands are expected to be associated with adverse health outcomes, it is less so for low job demands. In terms of *sleep regularity*, it may be that workers with low job demands have work schedules that are disruptive to regular sleep schedules. In contrast, workers with too many job demands may have more irregular sleep schedules driven by extensive work hours during the week. On the weekends, these workers with high job demands may need to extend their weekend awakenings due to the need for recovery from the workweek.⁴¹ For *sleep efficiency*, the

time needed to fall asleep at night could be influenced by how much restoration is needed and whether rumination is occurring. For example, having too few job demands may reflect a low intellectual engagement at work and limit the need for the restorative role of sleep⁴² thereby lowering the propensity for sleep. On the other hand, higher job demands have been found to relate to worse sleep quality due to rumination.⁴³ Sleep efficiency may be affected if an individual is unable to disconnect from work and engages in rumination because of their job demands.

We also found that the relationship between job demands and sleep health depended on the sleep dimension of focus. The most consistent linear relationships with overall and specific job demands indicators were found for sleep irregularity and sleep dissatisfaction. Regardless of the type of job demands, higher demands were related to more irregular and dissatisfied sleep. For the other sleep health dimensions (sleep duration, nap frequency, sleep inefficiency), effects of job demands were limited to certain types of demands, but similar linear relationships appeared. The one exception was the association between job intensity and nap frequency, which exhibited a negative linear relationship. It may be that workers with more intense work are too engaged in their work or too busy to take naps resulting in greater alertness during the day.

For stratified analyses, we found evidence suggesting that job control was a buffer for job demands, with results appearing mainly for participants with low job control. Thus, the negative effects of job demands on sleep health outcomes seem to be attenuated by higher job resources. One exception was sleep dissatisfaction—individuals with higher demands report more dissatisfied sleep regardless of their level of job control. Some evidence suggests that job demands, but not factors related to job resources, were related to future sleep problems.²¹ Others have also found that higher job demands were directly associated with poorer sleep quality, whereas higher job control was indirectly associated with better sleep quality through increased job satisfaction and reduced chronic fatigue.²⁰ It could be that job demands have a strong negative linear relationship with sleep quality or satisfaction regardless of job control, possibly due to the cumulative effects of this stressor exceeding individuals' stress resilience capacity.¹¹

Taken together, an increase in job demands—even to a moderate level—may not allow satisfying/restorative sleep in workers. However, for other sleep dimensions, quadratic effects were found more prevalently for individuals with low job control, with sleep inefficiency as the only sleep health dimension exhibiting a significant quadratic effect for participants with high job control. Overall, results revealed that sleep health in workers was optimal when the levels of job demands were moderate, and this was particularly so for those with low job control. This novel finding extends previous literature on work stress and sleep.

Strengths, limitations, and future directions

Our study has several strengths. We used a national sample of adults, which offered the opportunity to test our new hypotheses in a large and general sample of full-time workers. The assessment of overall demands and specific aspects of job demands (ie, intensity, role conflict, work overload, time pressure, and interruptions) and comparison of results across the 6 job demands indicators in relation to sleep outcomes are also strengths. We included 5 dimensions of sleep health, based on a well-validated conceptual model (RuSATED²⁴), which extends past work examining single sleep measures (eg, sleep duration or sleep quality only).

There are also some limitations. First, although we focused on job demands and job control given their potentially salient effects on employee health,^{15,16} there may be other work characteristics not captured by the job demands-control model that we did not include

but may be important for sleep health, such as perceived inequality at work,⁶ shift work,^{5,17} work precarity,^{44,45} salary/income (eg, experiences of poverty⁴⁶), or commute time.⁴⁷ Moreover, as the context for work is increasingly shifting to remote work arrangements, there is the need to understand how the context of work (ie, office environment vs. home environment) may relate to sleep health.⁴⁸ Additionally, there may be non-work-related factors that workers have to balance with their workloads such as demands from family/personal life⁴⁹ or caregiving⁵⁰ that may also influence sleep health. Although beyond the scope of the current study, future research would benefit from considering both (1) other work-related and (2) non-work-related characteristics when assessing sleep health outcomes to have a more nuanced understanding of the relationships among various factors that may relate to sleep.

Second, the cross-sectional design prevents examination of causal relationships between job demands and sleep. Inclusion of multiple time points would allow assessment of directionality of effects between job demands and sleep health. Third, we used self-reported measures of job demands and sleep health, which might have inflated the relationship due to the common self-report method.⁵¹ Including objective measures of job demands and control (eg, peer or supervisor ratings) and sleep (eg, actigraphy) could expand our understanding of the relationships between these variables.

Fourth, some components of the Ru-SATED²⁴ model (ie, sleep timing and WASO for sleep inefficiency) were not available in the MIDUS core survey and were not included. Fifth, the MIDUS sample was mostly non-Hispanic White; therefore, we do not know whether our findings would replicate in racially and ethnically diverse samples. Future research could assess whether patterns are similar across races/ethnicities and different age groups. Future research could also consider other potential moderators. For example, job tenure (a proxy for exposure to specific working conditions) may moderate the associations between job demands, job control, and sleep, but we did not control for this variable in analyses. Cumulative exposure to the stress of job demands could lead to either (1) loss spirals or (2) job crafting⁸ which could intensify or attenuate the effect of job demands on health outcomes, respectively. Further, cumulative exposure to chronic stressors is related to negative health outcomes.¹¹ Examining moderations by specific job conditions and demographic characteristics would be fruitful directions for future research.

Conclusions

To ensure the good health of workers, it is important to determine the most ideal levels of demands and control needed for the best working conditions and subsequent health outcomes that may be driven by sleep health. We found that moderate levels of job demands were related to optimal sleep health, especially for workers with low job control. Tailoring the work environment by assigning not too high and not too low job demands can assure that the workplace provides a fitting level of responsibilities that benefit employee sleep health. Alternatively, if extensive levels of job demands cannot be reduced to a moderate level, enhancing employees' job control in these situations may be another way to promote sleep health in workers.

Declarations of interest

None.

Funding

This work was supported by the National Institute on Aging under Grant: [R56AG065251](https://doi.org/10.1016/j.sleh.2022.09.002) to Soomi Lee (PI). The National Institute on Aging was not involved in the design or conduct of the study; data

management, analyses, or interpretation; or preparation, review, or submission of the manuscript for publication.

Acknowledgments

Disclosure Statement: Outside of the current work, Orfeu M. Buxton discloses that he received subcontract grants to Penn State from Proactive Life (formerly Mobile Sleep Technologies), doing business as SleepSpace (National Science Foundation grant #1622766 and NIH/National Institute on Aging Small Business Innovation Research Program [R43AG056250](https://doi.org/10.1016/j.sleh.2022.09.002), [R44 AG056250](https://doi.org/10.1016/j.sleh.2022.09.002)); received honoraria/travel support for lectures from Boston University, Boston College, Tufts School of Dental Medicine, University of Miami, Eric H. Angle Society of Orthodontists, Harvard Chan School of Public Health, New York University, and Allstate; receives consulting fees from SleepNumber; and receives an honorarium for his role as the Editor in Chief of Sleep Health [sleephealthjournal.org](https://doi.org/10.1016/j.sleh.2022.09.002).

Supplementary materials

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.sleh.2022.09.002](https://doi.org/10.1016/j.sleh.2022.09.002).

References

- 1 Kwok CS, Kontopantelis E, Kuligowski G, et al. Self-reported sleep duration and quality and cardiovascular disease and mortality: a dose-response meta-analysis. *J Am Heart Assoc*. 2018;7(15):1–26.
- 2 Sabia S, Fayosse A, Dumurgier J, et al. Association of sleep duration in middle and old age with incidence of dementia. *Nat Commun*. 2021;12(1):1–10.
- 3 Grandner MA. Sleep, health, and society. *Sleep Med Clin*. 2020;15(2):319–340.
- 4 Bannai A, Tamakoshi A. The association between long working hours and health: a systematic review of epidemiological evidence. *Scand J Work Environ Health*. 2014;40(1):5–18.
- 5 Jehan S, Zizi F, Pandi-Perumal SR, et al. Shift work and sleep: medical implications and management. *Sleep Med Disord Int J*. 2017;1(2):36–42.
- 6 Lee S, Mogle JA, Jackson CL, Buxton OM. What's not fair about work keeps me up: perceived unfairness about work impairs sleep through negative work-to-family spillover. *Soc Sci Res*. 2019;81:23–31.
- 7 Lawton MP. Environment and other determinants of well-being in older people. *Gerontologist*. 1983;23(4):349–357.
- 8 Bakker AB, Demerouti E. Job demands-resources theory: taking stock and looking forward. *J Occup Health Psychol*. 2017;22(3):273–285.
- 9 Reijseger G, Schaufeli WB, Peeters MCW, Taris TW, van Beek I, Ouweneel E. Watching the paint dry at work: psychometric examination of the Dutch Boredom Scale. *Anxiety Stress Coping*. 2013;26(5):508–525.
- 10 French KA, Allen TD, Henderson TG. Challenge and hindrance stressors in relation to sleep. *Soc Sci Res*. 2018;2019:145–153. 222.
- 11 Epel ES. The geroscience agenda: toxic stress, hormetic stress, and the rate of aging. *Ageing Res Rev*. 2020;63: 101167.
- 12 Yerkes RM, Dodson JD. The relation of strength of stimulus to rapidity of habit-formation. *J Comp Neurol Psychol*. 1908;18:459–482.
- 13 Calabrese EJ. Converging concepts: adaptive response, preconditioning, and the Yerkes-Dodson Law are manifestations of hormesis. *Ageing Res Rev*. 2008;7(1):8–20.
- 14 Karasek RA, Theorell T. *Healthy Work: Stress, Productivity and the Reconstruction of Working Life*. New York: Basic Books; 1990.
- 15 Xanthopoulou D, Bakker AB, Dollard MF, et al. When do job demands particularly predict burnout?: the moderating role of job resources. *J Manag Psychol*. 2007;22(8):766–786.
- 16 Litwiller B, Snyder LA, Taylor WD, Steele LM. The relationship between sleep and work: a meta-analysis. *J Appl Psychol*. 2017;102(4):682–699.
- 17 Linton SJ, Kecklund G, Franklin KA, et al. The effect of the work environment on future sleep disturbances: a systematic review. *Sleep Med Rev*. 2015;23:10–19.
- 18 Van Laethem M, Beckers DGJ, Kompier MAJ, Dijksterhuis A, Geurts SAE. Psychosocial work characteristics and sleep quality: a systematic review of longitudinal and intervention research. *Scand J Work Environ Heal*. 2013;39(6):535–549.
- 19 Garefelt J, Platts LG, Hyde M, Magnusson Hanson LL, Westerlund H, Akerstedt T. Reciprocal relations between work stress and insomnia symptoms: a prospective study. *J Sleep Res*. 2020;29(2):1–11.
- 20 Li Y, Fang J, Zhou C. Work-related predictors of sleep quality in Chinese nurses: testing a path analysis model. *J Nurs Res*. 2019;27(5):1–10.
- 21 Muhamad Nasharudin NA, Idris MA, Young LM. The effect of job demands on health and work outcomes: a longitudinal study among Malaysian employees. *PsyCh J*. 2020;9(5):691–706. <https://doi.org/10.1002/pchj.378>.

- 22 Stiglbauer B, Kovacs C. The more, the better? Curvilinear effects of job autonomy on well-being from vitamin model and PE-fit theory perspectives. *J Occup Health Psychol.* 2018;23(4):520–536.
- 23 Gadinger MC, Fischer JE, Schneider S, Fischer GC, Frank G, Kromm W. Female executives are particularly prone to the sleep-disturbing effect of isolated high-strain jobs: a cross-sectional study in German-speaking executives. *J Sleep Res.* 2009;18(2):229–237.
- 24 Buysse DJ. Sleep health: can we define it? Does it matter? *Sleep.* 2014;37(1):9–17.
- 25 Ryff CD, Krueger RF. Approaching Human Health as an Integrative Challenge: Introduction and Overview. In: Ryff CD, Krueger RF, eds. *The Oxford Handbook of Integrative Health Science.* New York: Oxford University Press; 2018:3–22.
- 26 Dijk DJ, Beersma DGM, Daan S. EEG power density during nap sleep: reflection of an hourglass measuring the duration of prior wakefulness. *J Biol Rhythms.* 1987;2(3):207–219.
- 27 Mantua J, Spencer RMC. Exploring the nap paradox: are mid-day sleep bouts a friend or foe? *Sleep Med.* 2017;37:88–97.
- 28 Cousins JN, Leong RLF, Jamaluddin SA, Ng ASC, Ong JL, Chee MWL. Splitting sleep between the night and a daytime nap reduces homeostatic sleep pressure and enhances long-term memory. *Sci Rep.* 2021;11(1):1–15.
- 29 Milner CE, Cote KA. Benefits of napping in healthy adults: impact of nap length, time of day, age, and experience with napping. *J Sleep Res.* 2009;18(2):272–281.
- 30 Pan Z, Huang M, Huang J, Yao Z, Lin Z. Association of napping and all-cause mortality and incident cardiovascular diseases: a dose–response meta analysis of cohort studies. *Sleep Med.* 2020;74:165–172.
- 31 Lee S, Smith CE, Wallace ML, et al. Cardiovascular risks and sociodemographic correlates of multidimensional sleep phenotypes in two samples of US adults. *SLEEP Adv.* 2022;3(1):1–11.
- 32 Li P, Gao L, Yu L, et al. Daytime napping and Alzheimer's dementia: a potential bidirectional relationship. *Alzheimer's Dement.* 2021;2022:1–11.
- 33 Dalmaes M, Benítez ID, Mas A, et al. Assessing sleep health in a European population: results of the catalan health survey 2015. *PLoS One.* 2018;13(4):1–13.
- 34 Lee S, Mu CX, Wallace ML, et al. Sleep health composites are associated with the risk of heart disease across sex and race. *Sci Rep.* 2022;12(1):1–11.
- 35 Karasek RA, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. The Job Content Questionnaire (JCQ): an instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol.* 1998;3(4):322–355.
- 36 Andel R, Crowe M, Kareholt I, Wastesson J, Parker MG. Indicators of job strain at midlife and cognitive functioning in advanced old age. *J Gerontol B Psychol Sci Soc Sci.* 2011;66(3):287–291.
- 37 Hawkins MAW, Gathright EC, Gunstad J, et al. The MoCA and MMSE as screeners for cognitive impairment in a heart failure population: a study with comprehensive neuropsychological testing. *Hear Lung J Acute Crit Care.* 2014;43(5):462–468.
- 38 Hickman RJ, Khambaty T, Stewart JC. C-reactive protein is elevated in atypical but not nonatypical depression: data from the National Health and Nutrition Examination Survey (NHANES) 1999–2004. *J Behav Med.* 2014;37(4):621–629.
- 39 Kline RB. *Principles and Practice of Structural Equation Modeling.* 2nd ed. New York: Guilford Press; 2005.
- 40 Parker KN, Ragsdale JM. Effects of distress and eustress on changes in fatigue from waking to working. *Appl Psychol Heal Well-Being.* 2015;7(3):293–315.
- 41 Myllyntausta S, Kronholm E, Pulakka A, et al. Association of job strain with accelerometer-based sleep duration and timing of sleep among older employees. *J Sleep Res.* 2022;31(2):1–10.
- 42 Akerstedt T, Nilsson PM, Kecklund G. Sleep and recovery. *Res Occup Stress Well Being.* 2009;2009(7):205–247.
- 43 Van Laethem M, Beckers DGJ, Geurts SAE, Garefelt J, Magnusson Hanson LL, Leineweber C. Perseverative cognition as an explanatory mechanism in the relation between job demands and sleep quality. *Int J Behav Med.* 2018;25(2):231–242.
- 44 Mai QD, Jacobs AW, Schieman S. Precarious sleep? Nonstandard work, gender, and sleep disturbance in 31 European countries. *Soc Sci Med.* 2019;237: 112424.
- 45 Salas-Nicás S, Sembajwe G, Navarro A, Moncada S, Llorens C, Buxton OM. Job insecurity, economic hardship, and sleep problems in a national sample of salaried workers in Spain. *Sleep Heal.* 2020;6(3):262–269.
- 46 Patel NP, Grandner MA, Xie D, Branas CC, Gooneratne N. “Sleep disparity” in the population: poor sleep quality is strongly associated with poverty and ethnicity. *BMC Public Health.* 2010;10:1–11.
- 47 Kim S, Kim Y, Lim SS, Ryoo JH, Yoon JH. Long commute time and sleep problems with gender difference in work life balance: a cross-sectional study of more than 25,000 workers. *Saf Health Work.* 2019;10(4):470–475.
- 48 Allen TD, Golden TD, Shockley KM. How effective is telecommuting? Assessing the status of our scientific findings. *Psychol Sci Public Interes.* 2015;16(2):40–68.
- 49 Buxton OM, Lee S, Beverly C, et al. Work-family conflict and employee sleep: evidence from IT workers in the work, family and health study. *Sleep.* 2016;39(10):1871–1882.
- 50 Gao C, Chapagain NY, Scullin MK. Sleep duration and sleep quality in caregivers of patients with dementia a systematic review and meta-analysis. *JAMA Netw Open.* 2019;2(8):1–15.
- 51 Podsakoff PM, MacKenzie SB, Lee JY, Podsakoff NP. Common method biases in behavioral research: a critical review of the literature and recommended remedies. *J Appl Psychol.* 2003;88(5):879–903.